

- 1. An isolated Robo polypeptide comprising SEQ ID NO:2, 4, 6, 8, 10 or 12, or a polypeptide domain thereof having at least 12 consecutive residues thereof and a Robospecific activity, wherein said domain is encoded by neither EST yq76e12 nor yq76e12.
- 2. An isolated polypeptide according to claim 1, wherein said activity is selected from at least one of a Robo-competitive binding, Robo-specific antigenicity and a Robo-specific immunogenicity.
- 3. An isolated polypeptide according to claim 1, wherein said domain comprises at least one of a Robo immunoglobulin, fibronectin or cytoplasmic motif domain.
- 4. A recombinant nucleic acid encoding a polypeptide according to claim 1.
- 5. A cell comprising a nucleic acid according to claim 4.
- 6. A method of making a Robo polypeptide, comprising the following steps: incubating a host cell or cellular extract containing a nucleic acid according to claim 4 under conditions whereby the polypeptide encoded by the nucleic acid is expressed and recovering the expressed polypeptide.
- 7. An isolated Robo polypeptide made by the method of claim 6.
- 8. An isolated *robo* nucleic acid comprising a strand of SEQ ID NO:1, 3, 5, 7, 9 or 11, or a fragment thereof having at least 24 consecutive bases thereof, and sufficient to specifically hybridize with a nucleic acid having the sequence defined by the corresponding opposite strand, wherein the fragment is contained in neither EST yq76e12 nor yq76e12.
- 9. A method for modulating cell function or morphology comprising providing the cell with an agent which modulates activity of a Robo polypeptide or function of a *robo* gene, wherein the agent comprises a polypeptide according to claim 1 or a Robo-specific antibody.

Add A' add Die >